

PROFESSOR ECKART HANEKE (Orcid ID : 0000-0001-9957-1441)

Article type : Original Article

***Fusarium* onychomycoses in Switzerland – a mycological and histopathological study**

Anna Rammlmair¹, Konrad Mühlethaler,² Eckart Haneke^{1,3,4,5}*¹

¹Dept Dermatol, Inselspital, Univ Berne, Bern, Switzerland

² Inst Infectious Diseases, Univ Berne, Bern, Switzerland

³ Dermatol Practice Dermaticum, Freiburg, Germany

⁴ Centro Dermatol, Inst CUF, Matosinhos, Porto, Portugal

⁵ Kliniek Huidziekten, Univ Hosp, Gent, Belgium

The authors declare that there is no conflict of interest.

Summary

Objectives: Onychomycoses in temperate climates are most commonly due to dermatophytes, particularly *Trichophyton rubrum*. Non-dermatophyte nail infections are much less frequent, and their diagnosis requires a careful and repeated search for a potential dermatophyte that may have been overgrown in culture.

Materials/Patients/Methods: A series of histological slides of suspected onychomycoses with uncommon fungal morphology prompted us to search for non-dermatophytic molds causing dermatophytosis-like nail infections.

* Corresponding author: Prof Dr Eckart Haneke, Schlippehof 5, 79110 Freiburg, Germany. Email haneke@gmx.net, Tel +49 761 8978368, Fax +49 761 3837401

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/myc.12964

This article is protected by copyright. All rights reserved.

Results and Conclusions: Thirty cases were identified by culture as *F solani*, *F oxysporum*, *F dimerum* or *F spp*, and two more were only diagnosed histopathologically. None of these patients was immunocompromised. Treatment was mostly unsuccessful with terbinafine whereas itraconazole showed a moderately better treatment result; in all cases, a topical ciclopirox nail varnish in a hydroxychitosan base was added. The study demonstrates that *Fusarium* species are important nail pathogens also in Central Europe.

Key words: Onychomycosis – *Fusarium* species – Culture – Histopathology – Treatment

Introduction

Onychomycoses (OM) are worldwide occurring fungal infections of the nail apparatus. Most cases are distal or distal-lateral subungual OM and due to dermatophytes [1]. Yeasts and other non-dermatophytic molds are rare and not generally accepted as nail pathogens; however, some *Candida* species have acid proteases capable of digesting keratin, and some molds, including *Fusarium* spp are undoubtedly able to invade nails and subungual keratin [2]. English (1979) stated that molds should only be accepted as nail pathogens when the same species is grown three times in culture and without simultaneous growth of a dermatophyte [3]. These stringent criteria are rarely fulfilled in clinical routine although they might be useful for the choice of an antifungal drug, particularly in recalcitrant cases [4].

Fusarium species are common plant pathogens and occur both in tropical as well as temperate and even arctic climates [5]. They cause important crop damages and are the source of mycotoxins also in Central Europe including Switzerland [6]. Of the more than 300 known phylogenetically different species, most opportunistic *Fusarium* pathogens belong to the *F solani* complex, *F oxysporum* complex and *F fujikoro*i complex. In tropical countries, *Fusarium* spp cause keratitis and onychomycoses. They are also feared causes of systemic infections in severely immunocompromised individuals, particularly under chemotherapy and in acute leukemias [7]. Onychomycoses in immunocompetent persons due to *Fusarium* spp in Europe are mainly reported from southern countries

such as the Balkans, Italy, France and Spain and mostly occur in toes that often have been traumatized before [8, 9]. They appear to be more frequent in women than in men [10, 11].

During our routine examinations of onychomycosis-suspicious nails using direct microscopy, culture and histopathology, we encountered a number of cases with an uncommon fungal morphology (Figure 1) prompting us to search for the cause.

Material and methods

The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines page, have been adhered to. No ethical approval was required as the research in this article related to micro-organisms.

The culture results of nail specimens from 2010 till 2016 were searched from the files of the Mycology laboratory, Institute for Infectious Diseases, Univ Berne and compared with the results of the histological examinations. As direct microscopy of KOH cleared specimens was noncontributory and usually did not give a hint at a particular *Fusarium* etiology, the presence of fungal elements was noted but not included into the final analysis. Cultures were grown on Sabouraud agar with gentamycin and chloramphenicol, Czapek's agar, mycosel agar and the micromorphology was evaluated using lacto fuchsin stain. Histological sections were prepared from nail clippings with as much subungual keratin as possible, stained with hematoxylin and eosin (H&E) and PAS and viewed at 100, 200 and 400x magnifications. In case of doubt, Grocott silver stain was performed.

Results

The institutional files revealed 28 cases of *Fusarium* spp, of which 15 were *F solani*, 5 were *F oxysporum*, 7 were *F* species, and 1 was *F dimerum* (Figure 2). Two more cases were seen in Freiburg, Germany, and another two in Porto, Portugal (EH^{3,4}); of these four, two were mycologically diagnosed as *F* species and 2 were only diagnosed histopathologically.

Histopathology of the clipped nail and subungual keratosis demonstrated fungal elements in the deep nail plate layer and the underlying keratosis. Leukocyte accumulations as seen in Munro's microabscesses were common and surrounded by parakeratosis. PAS exhibited fungal filaments with very variable diameter and often

ampullar dilatations. The hyphae showed marked segmentation with frequently very short segments (Figure 1). Nine-teen of the specimens demonstrated basophilic fungal cell walls and thus allowed the diagnosis of fungal infection also in H&E sections (Figure 1a).

Discussion

Fusarium species are new emergent nail pathogens now observed almost worldwide. Originally known as plant pathogens in warm climate zones with an important potential to harm crops and as a source of fungal mycotoxins in cereals they were observed as causes of *Fusarium* keratitis in rural areas mainly, but not exclusively, of tropical countries and as occasional pathogens of onychomycoses in tropical and subtropical regions. *Fusarium* nail infections are rarely diagnosed in Central Europe and difficult to treat. Clinically, they cannot be distinguished from the common dermatophyte onychomycoses, but histologically they show slender to relatively thick septate hyphae often with characteristic ampullar dilatations in their course. In culture, they form whitish cottony colonies with a brownish reverse side. Lactophenol cotton blue stain reveals characteristic banana shaped macroconidia in addition to microconidia and hyphae.

Nondermatophyte molds cause about 10% of the onychomycoses with a range from 1.45% to 17.6% [12]. *Fusarium* spp alone were estimated at 1 to 6% of onychomycoses [13], but were found in a higher percentage in other studies [14, 15, 16]. The differences in prevalence may account to various geographical regions and climate zones and certainly also to techniques of examination [17]. It is evident that *Fusarium* onychomycoses go unnoticed in most cases and are often simply dismissed as another case of therapy-resistant onychomycosis or as mold contamination. The percentage of *Fusarium* spp as nail pathogens varies from 9 – 44% of mold infections [18]. Women are more frequently affected and fingernails are apparently also involved in tropical countries [16]. All types of onychomycoses may be due to *Fusarium* spp [9, 19] including onychomycoses with associated paronychia [20] and melanonychia [21]. Whether diabetes mellitus is a risk factor is not yet entirely clear [22].

Fusarium onychomycoses are important as they may be the source of life-threatening infections in immunocompromised individuals and cause deep seated fusarioses in burns, large ulcers and foreign body granulomas [23, 24]. Patients with these infections should not work in the food industry, kitchens and hospitals, particularly newborn and cancer wards as well as surgical theaters.

Treatment of *Fusarium* infections is difficult. Fluconazole is usually inactive and also terbinafine often only gives a modest response. Itraconazole is more active [25], and both voriconazole and high-dose amphotericine are currently the drugs of choice [26, 27]. Natamycin was found to be very active against *F. fujikuroi* [28]. One case was reportedly cured with 2 sessions of Nd-YAG laser [29] whereas another study showed no cure with laser treatment [30].

In summary, *Fusarium* species are emergent pathogens also in Central Europe. They should be searched for in case of mold growth in culture and an uncommon morphology in histopathology of the nail. Their identification is important as they are more difficult to treat and may have particular hygienic implications for a variety of professions and contact with premature babies, immunosuppressed individuals, and cancer patients.

References

1. Baran R, Hay RJ, Tosti A, Haneke E. A new classification of onychomycoses. Br J Dermatol 1998;139:567-571
2. Galletti J, Negri M, Grassi FL, Kioshima-Cotica ÉS, Svidzinski TI. *Fusarium* spp. is able to grow and invade healthy human nails as a single source of nutrients. Eur J Clin Microbiol Infect Dis. 2015;34:1767-1772
3. English MP. Nails and fungi. Br J Dermatol 1976;94:697-701
4. Tupaki-Sreepurna A, Jishnu BT, Thanneru V, Sharma S, Gopi A, Sundaram M, Jyoti Kindo A. An assessment of in vitro antifungal activities of efinaconazole and itraconazole against common non-dermatophyte fungi causing onychomycosis. J Fungi (Basel) 2017;3(2). pii: E20. doi: 10.3390/jof3020020
5. Hocquette A, Grondin M, Bertlout S, Mallié M. Les champignons des genres *Acremonium*, *Beauveria*, *Chrysosporium*, *Fusarium*, *Onychocola*, *Paecilomycetes*, *Penicillium*, *Scedosporium* et *Scopulariopsis* responsables de hyalohyphomycoses. J Mycol Méd 2005;15:136-149

- Accepted Article
6. Vogelgsang S, Musa T, Bänziger I, Kägi A, Bucheli TD, Wettstein FE, Pasquali M, Forrer HR. Fusarium mycotoxins in Swiss wheat: A survey of growers' samples between 2007 and 2014 shows strong year and minor geographic effects. *Toxins (Basel)* 2017;9(8). pii: E246. doi: 10.3390/toxins9080246
 7. Haneke E. Mucor mycosis and neonatal fat necrosis in a premature neonate. Clear-cell syringomatous carcinoma. Fusariosis. *Coll Dermato-Pathol Unna-Darier, Zürich*, 1. – 3. 9. 2000
 8. Nucci M, Varon AG, Garnica M, Akiti T, Barreiros G, Trope BM, Nouér SA. Increased incidence of invasive fusariosis with cutaneous portal of entry, Brazil. *Emerg Infect Dis* 2013;19:1567-1572
 9. Baran R, Tosti A, Piraccini BM. Uncommon clinical patterns of Fusarium nail infection: report of three cases. *Br J Dermatol* 1997;136:424-427
 10. Calado NB, Sousa F jr, Gomes NO, Cardoso FR, Zaror LC, Milan EP. Fusarium skin and nail infection: A report of eight cases from Natal, Brazil. *Mycopathologia* 2006;161:27-31
 11. Yang YS, Ahn J-J, Shin M-K, Lee M-H. Fusarium solani onychomycosis of the thumbnail coinfecting with Pseudomonas aeruginosa. Report of two cases. *Mycoses* 2011;54:168-171
 12. Bao F, Fan Y, Sun L, Yu Y, Wang Z, Pan Q, Yu C, Liu H, Zhang F. Comparison of fungal fluorescent staining and ITS rDNA PCR-based sequencing with conventional methods for the diagnosis of onychomycosis. *J Eur Acad Dermatol Venereol* 2018;32:1017-1021
 13. Guilhermetti E, Takahachi G, Shinobu CS, Svidzinski TI. Fusarium spp. as agents of onychomycosis in immunocompetent hosts. *Int J Dermatol* 2007;46:822-826
 14. Lawry MA, Haneke E, Strobeck K, Martin S, Zimmer B, Romano PS. Methods for diagnosing onychomycosis: a comparative study and review of the literature. *Arch Dermatol* 2000;136:1112-1116
 15. Zukervar P, Dabin G, Secchi T, Petiot-Roland A, Mathon N, Maccari M, Pincemaille B, Colombet-Navaranne A, Rigot-Muller G, Batut V, Picot S, Bienvenue AL. Etude des onychomycoses en médecine de ville dans la région lyonnaise. *J Mycol Méd* 2011;21:118-122
 16. Diongue K, Ndiaye M, Seck MC, Diallo MA, Badiane AS, Ndiaye D. Onychomycosis caused by *Fusarium* spp. in Dakar, Senegal: Epidemiological,

clinical, and mycological study. *Dermatol Res Pract* 2017;2017:1268130. doi: 10.1155/2017/1268130 Epub 2017 Dec 4


17. Nelson PE, Digniani MC, Anaissie EJ. Taxonomy, biology and clinical aspects of *Fusarium* species. *Clin Microbiol Rev* 1994;7:479-504
18. Kaufmann-Lacroix C, Villers A, Gantier JC, Guillet G, Wierzbicka E, Rodier MH. Onyxis and cutaneous ulcers due to *Fusarium solani* in a patient with diabetes mellitus. *J Mycol Méd* 2005;15:150-154
19. Cobos Llardó D, Fierro Arias L, Arellano Mendoza I, Bonifaz A. Onychomycosis and its influence on quality of life. *Dermatol Cosm Med Quir* 2016;14:318-327
20. Ranawaka RR, Nagahawatte A, Gunasekara TA. *Fusarium* onychomycosis: prevalence, clinical presentations, response to itraconazole and terbinafine pulse therapy, and 1-year follow-up in nine cases. *Int J Dermatol* 2015;54:1275-1282
21. Park KY, Lee JW, Kim IS et al. Longitudinal melanonychia caused by *Fusarium oxysporum* in immunocompetent patient. *Mycoses* 2012;55:e40-41
22. Wu CY Chen GS, Lan CC. Onychomycosis caused by *Fusarium solani* in a woman with diabetes. *Clin Exp Dermatol* 2009;34:e772-774
23. Bourgeois GP, Cafardi JA, Sellheyer K, Andea AA. Disseminated *Fusarium* infection originating from paronychia in a neutropenic patient: a case report and review of the literature. *Cutis* 2010;85:191-194
24. Ocampo Garza J Herz-Ruelas ME, Chavez-Alvarez S, Gómez-Flores M, Vera-Cabrera L, Welsh-Lozano O, Gallardo-Rocha A, Escalante-Fuentes WG, Ocampo-Candiani J. Disseminated fusariosis with endophthalmitis after skin trauma in acute lymphoblastic leukaemia. *J Eur Acad Dermatol Venereol* 2016;30:e121-e123
25. Gupta AK, De Doncker P, Scher RK, Haneke E, Daniel CR 3rd, André J, Baran R. Itraconazole for the treatment of onychomycosis. *Int J Dermatol* 1998;37:303-308
26. Garbino J, Uckay I, Rohner P, Lew D, Van Delden C. *Fusarium* peritonitis concomitant to kidney transplantation successfully managed with voriconazole: case report and review of the literature. *Transpl Int* 2005 ;18:613-618


27. Khetan S, Khetan P, Katkar V, Kusulkar M. Urinary tract infection due to *Fusarium oxysporum* in an immunocompetent patient with chronic kidney disease. J Biomed Res. 2018 ;32:157-160
28. Tupaki-Sreepurna A, Thanneru V, Natarajan S, Sharma S, Gopi A, Sundaram M, Kindo AJ. Phylogenetic diversity and in vitro susceptibility profiles of human pathogenic members of the *Fusarium fujikuroi* species complex isolated from South India. Mycopathologia 2018;183:529-540
29. Khurana A, Chowdhary A, Sardana K, Gautam RK, Sharma PK. Complete cure of *Fusarium solani* sp. complex onychomycosis with Qs NdYAG treatment. Dermatol Ther 2018;31:e12580
30. Espírito-Santo GAd, Pereira Leite D, Hoffmann-Santos HD, Basili Dias L, Hahn RC. 1340 nm laser therapy for onychomycosis: Negative results of prospective treatment of 72 toenails and a literature review. J Clin Aesthet Dermatol 2017;10:56-61

Legends

Figure 1. Onychomycosis due to *Fusarium solani*. Both the hematoxylin & eosin (H&E) (1a) as well as the PAS stain (1b) show fungal filaments of variable diameter with some ampullar dilatations. Usually, H&E does not stain fungi, particularly not dermatophytes, The fungi invade the deep nail plate layer as well as the parakeratosis surrounding a Munro-abscess like accumulation of neutrophils. Original magnification x400.

Figure 2. Distribution of culturally identified *Fusarium* species.





28 cases of Fusarium spp.

